

Cal/Ecotox
Exposure Factors for American Kestrel (Falco sparverius)*

Page 1

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Age at Fledging, Metamorphosis, Weaning			28-30	d	B	Fledgling	Lassen; CA	a	1
Age at Fledging, Metamorphosis, Weaning	28.4		26-32	d	B	Fledgling	Lab	b	2
Age at Fledging, Metamorphosis, Weaning			29-31	d	B	Fledgling	Nevada; Sierra; CA	c	3
Body Fat (total or %)	2.46	0.25 SE	0.8-4.9	g	B	Adult	Lab	d	4
Body Fat (total or %)	8.51	2.00 SE		%	F	Adult	NJ	e	5
Body Fat (total or %)			4-5.3	%	M	Adult	UT	f	6
Body Fat (total or %)	6.01	1.92 SE		%	M	Adult	CO; NJ	g	5
Body Weight - Mean	119.0		86.0-164.8	g	F	Adult	USA	h	7
Body Weight - Mean	120.2	5.3 SE		g	F	Adult	Lab	i	8
Body Weight - Mean	102.5		80.0-143.0	g	M	Adult	USA	j	7
Body Weight - Mean	113.4	2.0 SE		g	M	Adult	Lab	k	8
Body Weight - Mean	119.8	0.9 SE	117.4-123.9	g	F	Both Adult and Juv.	CA	l	9
Body Weight - Mean	110.7	1.1 SE	105.6-112.1	g	M	Both Adult and Juv.	CA	m	9
Body Weight - Mean			10-12	g	B	Hatchling	Nevada; Sierra; CA	n	3
Clutch or Litter Size	4.3		3.5-4.9	eggs/clutch	F	Adult	Lassen; CA	o	1
Clutch or Litter Size	4.59	0.60 SD		eggs/clutch	F	Adult	CA	p	10
Clutch or Litter Size	5.1		4.8-5.2	eggs per nest	F	Adult	MO	q	11
Clutch or Litter Size	4			eggs/clutch	F	Adult	Nevada; Sierra; CA	r	3
Clutch or Litter Size	4.23		1-6	eggs	F	Adult	PA	s	12
Clutch or Litter Size	3.66		3-4	eggs	F	Adult	Lab	t	13
Clutch or Litter Size	4.4		5 (max)	eggs/clutch	F	Adult	WY	u	14
Clutch or Litter Size	4.4			eggs/clutch	F	Adult	IA	v	15
Clutch or Litter Size	4.1			young/nest	NR	Juvenile	WI	w	16
Dietary Composition	insects (31.7%), reptiles (26.0%), birds (16.6%), mammals (25.7%)				B	Adult	Nevada; Sierra; CA	x	3
Dietary Composition	mammal (82%), insect (16%), other (2%)				B	Adult	Yolo; CA	y	17
Dietary Composition	mammal (57%), insect (42%), other (1%)				B	Adult	Yolo; CA	z	17
Dietary Composition	mammal (24%), insect (74%), other (2%)				B	Adult	Yolo; CA	aa	17
Dietary Composition	mammal (43%), insect (51%), other (6%)				B	Adult	Yolo; CA	ab	17
Dietary Composition	mammal (39%), insect (60%), other (6%)				B	Adult	Yolo; CA	ac	17
Dietary Composition	invertebrates (173.8), reptiles (23.0), birds (481.0), mammals (609.5)				g	F	San Benito; Santa Clara; CA	ad	18
Dietary Composition	invertebrates (157.5), reptiles (0), birds (272.4), mammals (221.8)				g	M	San Benito; Santa Clara; CA	ae	18
Dietary Composition	invertebrates (39.3%), herpetofauna (37.9%), mammals (22.8%)				F	NR	Humboldt; CA	af	19
Dietary Composition	invertebrates (10.5%), herpetofauna (2.4%), mammals (79.1%), birds (8.0%)				F	NR	Humboldt; CA	ag	19

Exposure Factors for American Kestrel (Falco sparverius)

Page 2

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Dietary Composition	invertebrates (85.4%), vertebrates (14.6%)				NR	NR	Humboldt; CA	ah	20
Dietary Composition	insects (97%), isopod (5%), birds (27%), mammals (25%)				NR	NR	Tulare; CA	ai	21
Dietary Composition	white-footed mice (6.0%), meadow mice (78.9%), shrews (1.3%), pocket gopher (1.0%), uinta ground squirrel (0.3%), sm-med sized birds (5.7%), insects (59.9%)			%	NR	NR	WY	aj	14
Duration of Incubation or Gestation	27	0.58 SE	26-28	d	B	Embryo	Lab	ak	2
Duration of Incubation or Gestation			29-31	d	B	Embryo	Nevada; Sierra; CA	al	3
Duration of Incubation or Gestation	30.9		28-35	d	B	Embryo	PA	am	12
Duration of Incubation or Gestation	28.4		27-33	d	B	Embryo	Lab	an	13
Fledging or Weaning Rate	49%		42-57	%	B	Adult	PA	ao	22
Fledging or Weaning Rate	3.1		2.5-3.6	fledged/active nest	B	Fledgling	Lassen; CA	ap	1
Fledging or Weaning Rate	3.7		2.5-4.6	fledged/successful nest	B	Fledgling	Lassen; CA	aq	1
Fledging or Weaning Rate	5.0		3.5-5.46	number of fledglings per nest	B	Fledgling	MO	ar	11
Fledging or Weaning Rate	98			%	B	Fledgling	Nevada; Sierra; CA	as	3
Fledging or Weaning Rate	2.2			fledglings per nest	B	Fledgling	Sacramento; CA	at	23
Fledging or Weaning Rate	4.0	0.41		fledges/nest	NR	Fledgling	WA	au	24
Fledging or Weaning Rate	4.2			fledglings/pair	NR	Fledgling	WY	av	14
Fledging or Weaning Rate	88%			%	NR	Juvenile	WI	aw	16
Fledging or Weaning Rate	89%			%	NR	Juvenile	IA	ax	15
Food Ingestion Rate	0.42			kcal/g/day	M	Adult	Lab	ay	25
Food Ingestion Rate	43.6		28.0-70.2	cal/d	M	Adult	Lab	az	26
Food Ingestion Rate			18-21	g	B	NR	Lab	ba	14
Foraging Distance			<0.5-3.5	km	F	Adult	Yolo; CA	bb	27
Growth Rate	y = 7.77 + 7.44x				F	Nestling	Nevada; Sierra; CA	bc	3
Growth Rate	0.239				F	Nestling	Lab	bd	8
Growth Rate	y = 8.89 + 7.33x				M	Nestling	Nevada; Sierra; CA	be	3
Growth Rate	0.250				M	Nestling	Lab	bf	8
Hatching Success	79		70-83	%	B	Hatchling	Lassen; CA	bg	1
Hatching Success	89.3			%	B	Hatchling	Nevada; Sierra; CA	bh	3
Hatching Success	68%			%	NR	Juvenile	IA	bi	15
Hatching Success	2			%	NR	Nestling	WY	bj	14
Home Range	109.4			ha	B	Adult	Nevada; Sierra; CA	bk	3
Home Range	1.5			mi	NR	NR	IL	bl	28
Home Range	1.4			mi	NR	NR	IL	bm	28
Inhalation Rate	35.4	3.21 SD		#/min	NR	Adult	Lab	bn	29

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Longevity	12			yr	NR	Adult	Lab	bo	30
Longevity	11-07			yr-mo	NR	Adult	USA	bp	31
Longevity	118			mo	B	Both Adult and Juv.	USA	bq	32
Metabolic Rate			1.146-1.682	ml	B	Adult	Lab	br	33
				oxygen/g/hr					
Metabolic Rate	40.8			kcal/d	NR	Adult	Lab	bs	34
Metabolic Rate	23.06	1.03 SE	21.83-25.10	kcal/d	NR	Adult	Lab	bt	35
Population Density	23			individuals per 96 km2	B	Adult	Yolo; CA	bu	27
Population Density	50			individuals in 175 km2	B	Adult	MO	bv	36
Population Density			17-27	individuals per 52 km^2	B	Adult	OH	bw	37
Population Density	0.06		0.05-0.13	pairs per km2	B	Adult	MO	bx	11
Population Density	20			individuals/ 1476 sq km	B	Adult	WA	by	24
Population Density	10			pairs/1476 sq km	B	Adult	WA	bz	24
Population Density	1			pair/1.1 mi^2	B	Adult	WY	ca	14
Population Density	115	+/- 25 95% CI		individuals/ 100 mi	B	Both Adult and Juv.	Colusa; Humboldt; Lake; Mendocino; Yolo; CA	cb	38
Population Density	34	+/- 11 95% CI		kestrels/10 0 mi	B	Both Adult and Juv.	Colusa; Humboldt; Lake; Mendocino; Yolo; CA	cc	38
Population Density	48			individuals in 175 km2	B	Both Adult and Juv.	MO	cd	36
Population Density	0.22			individuals per km2	B	Both Adult and Juv.	MO	ce	11
Population Density	0.58			birds/mi2	NR	NR	IL	cf	28
Population Density	4			individuals/ 43 sq mi	NR	NR	IL	cg	28
Surface Area	7.15 x 10E-3			m^2	NR	Adult	Lab	ch	39
Surface Area	7.15 x 10E-3			m^2	NR	Adult	Lab	ci	39
Survival/ Mortality	0.465	0.017 SD			B	Adult	CANADA; USA	cj	10
Survival/ Mortality	0.602	0.035 SD			B	Both Adult and Juv.	CA	ck	10
Survival/ Mortality	12.6			mo	B	Both Adult and Juv.		cl	32
Survival/ Mortality	57			%	B	Both Adult and Juv.	USA	cm	7
Survival/ Mortality	78			%	B	Hatchling	PA	cn	12
Territory Size	1.4			km	B	Adult	OH	co	37
Territory Size	31.6	10.7 SD	18.7-42.0	ha	F	Adult	San Benito; Santa Clara; CA	cp	18
Territory Size	13.1	2.0 SD	9.7-14.8	ha	M	Adult	San Benito; Santa Clara; CA	cq	18
Territory Size			0.16-1.93	mi^2	NR	Adult	WY	cr	14
Territory Size			0.65-2.32	mi^2	NR	Adult	MI	cs	14

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Time of Fledging or Metamorphosis	July 29		July 5 - August 25		B	Fledgling	Lassen; CA	ct	1
Time of Hatching or Parturition	June 21		June 7- July 26		B	Hatchling	Lassen; CA	cu	1
Time of Hatching or Parturition	Jun 20-Jun 22				NR	Nestling	WY	cv	14
Time of Mating/ Laying	May 22		May 6 - June 26		F	Adult	Lassen; CA	cw	1
Time of Mating/ Laying	March-May				F	Adult	Lab	cx	2
Time of Mating/ Laying	late May - June				F	Adult	Nevada; Sierra; CA	cy	3
Time of Migration or Dispersal	August (begin)				B	Both Adult and Juv.	CA	cz	40
Time of Migration or Dispersal	February (peak)				B	Both Adult and Juv.	CA	da	40
Time of Migration or Dispersal	August (begin), October (peak), November (end)				B	NR	Marin; CA	db	41
Time of Molt	May - September				B	Adult	Nevada; Sierra; CA	dc	3
Time of Molt	July-October/November				B	Adult	USA	dd	7
Time of Nesting	May 20 to Jul 20				B	Adult	WY	de	14

- Notes**
- a N=112; Great Basin (elev. 1260-2340 m)
 - b mean nestling period; N=29 chicks; spring
 - c N=NR; Sagehen Creek Field Station
 - d total body fat based on Soxhlet fat extraction; N=21
 - e percent body fat calculated from total body electrical conductivity; N=13; Condition=migrating; Cape May
 - f mean percent body fat as measured by fat extraction from April to September; N=14; Condition=premigratory; April, July, September; Cache county
 - g percent body fat calculated from total body electrical conductivity; N=12; Condition=migrating; Cape May
 - h N=72; throughout USA
 - i average body weight; N=26 birds
 - j N=88; throughout USA
 - k average body weight; N=25 birds
 - l mean body weight measured over 4 years; N=111; coastal southern CA; see citation for body weights by region and season
 - m mean body weight measured over 4 years; N=69; coastal southern CA; see citation for body weights by region and season
 - n body weight at hatch; N=NR; Sagehen Creek Field Station
 - o N=38 clutches; Great Basin (1260-2340 m)
 - p N=244; unpublished data
 - q mean clutch size over 3 years; N=28 pairs; Condition=Breeding; rural and urban lands
 - r average clutch size; N=42; Sagehen Creek Field Station
 - s N=13 clutches; spring; Berks county
 - t mean clutch size in captivity; N=5; Condition=Breeding
 - u N=10 nests; spring; Jackson Hole
 - v mean clutch size (4 sample years); N=7-23 nests/year; Lucas and Wayne Counties (40deg57'N, 93deg18'W); All data are from artificial nest boxes.
 - w mean brood size (13 sample years); N=37 successful nests; Portage County (44deg27'N, 89deg40'W)
 - x percent of prey by prey weight; N=NR; Condition=breeding; Sagehen Creek Field Station; see citation for dietary composition by occurrence or cube-root of prey body weight
 - y percent composition of sample pellets (by volume); N=116 pellets; Jun.- Jul.; agricultural lands; values estimated from citation figure
 - z percent composition of sample pellets (by volume); N=59 pellets; Condition=breeding; Feb.-Mar.; agricultural lands; values estimated from citation figure
 - aa percent composition of sample pellets (by volume); N=104 pellets; Condition=breeding; May-Jun.; agricultural lands; values estimated from citation figure
 - ab percent composition of sample pellets (by volume); N=71 pellets; Condition=breeding; Mar.-Apr.; agricultural lands; values estimated from citation figure
 - ac percent composition of sample pellets (by volume); N=95 pellets; Condition=breeding; Apr.-May; agricultural lands; values estimated from citation figure
 - ad amount of prey taken by prey weight; N=5; winter; Hollister Basin (elev. 48-210 m); see citation for detailed dietary description
 - ae amount of prey taken by prey weight; N=5; winter; Hollister Basin (elev. 48-210 m); see citation for detailed diet description
 - af dietary composition (% of total biomass), 1973-1974; N=3; winter; Arcata Bottoms

Exposure Factors for American Kestrel (*Falco sparverius*)

Page 5

ag	dietary composition (% of total biomass), 1972-1973; N=4; winter; Arcata Bottoms
ah	relative numbers of dietary items captured; N=24 birds; October-February; Arcata Bottoms; pasture
ai	percent occurrence in sample pellets; N=NR; agricultural lands
aj	percent occurrence of prey items in pellets during nesting; N=299 pellets from 8 nests; spring; Jackson Hole
ak	mean number of days between 5th egg laid and 5th egg hatched; N=3 eggs; spring
al	N=NR; Sagehen Creek Field Station
am	incubation period; N=2 clutches (8 eggs); spring; Berks county
an	mean incubation period in captivity; N=5; Condition=Breeding
ao	mean percent of successful nests per year; N=259 nesting attempts; Berks and Lehigh counties; Successful nests were defined as those which fledged at least one young.
ap	N=36 nests; Great Basin (1260-2340 m)
aq	N=30 nests; Great Basin (1260-2340 m)
ar	three year mean fledging rate; N=28 pairs; Condition=Breeding; rural and urban lands
as	of chicks that hatched; N=150; Sagehen Creek Field Station
at	fledging rate per successful nest; N=17; spring; American River; oak woodlands, grasslands, residential
au	average number of young fledged; N=4 nests; Hanford Site, Benton and Franklin Counties
av	number of fledglings produced per pair; N=10 nests; spring; Jackson Hole
aw	percent of nests that fledged at least one young; N=42 nests; Portage County (44deg27'N, 89deg40'W)
ax	mean percent of young hatched that fledged (4 sample years); N=148 hatchlings; Lucas and Wayne Counties (40deg57'N, 93deg18'W); All data are from artificial nest boxes.
ay	mean caloric consumption per g body weight per day (wet weight basis); N=2; August
az	average and range of daily caloric intake (food provided ad libidum); N=1 bird; Condition=sedentary, injured
ba	averages of food amounts eaten per day; N=3 birds; Condition=captive; spring
bb	distance foraged from nest; N=4; Condition=Breeding; February 19-July 8; agricultural lands (lat, 38 deg 32'N; long., 121 deg 47'W)
bc	growth rate from day 3-16; N=NR; Sagehen Creek Field Station; see citation for growth curve
bd	growth rate of body weight in parent-reared birds; N=8 birds; See citation for growth rate of hand-reared birds.
be	growth rate from day 3-16; N=NR; Sagehen Creek Field Station; see citation for growth curve
bf	growth rate of body weight in parent-reared birds; N=11 birds; See citation for growth rate of hand-reared birds.
bg	mean hatchability measured over 4 years; N=157 boxes; Great Basin (elev. 1260-2340 m)
bh	N=168; Sagehen Creek Field Station
bi	mean percent of eggs hatched of total eggs laid (4 sample years); N=217 eggs; Lucas and Wayne Counties (40deg57'N, 93deg18'W); All data are from artificial nest boxes.
bj	percent of eggs that failed to hatch; N=10 nests; spring; Jackson Hole
bk	average home range - breeding; N=32; Sagehen Creek Field Station
bl	mean maximum diameter of winter home ranges; N=3 birds; winter; agricultural lands; based on distances between extreme observation points for individual kestrels
bm	mean maximum diameter of breeding home ranges; N=4 birds; spring; agricultural lands; based on distances between extreme observation points for individual kestrels
bn	mean resting respiratory rate; N=4; Condition=crippled, non-releasable; corresponding mean body weight was 110 g
bo	N=1; captive individual
bp	from USFWS Bird Banding Laboratory data; N=1832 band recoveries
bq	longest survival of a banded individual; N=1,017 records of mortality
br	combined day and night oxygen consumption over one 24 hr period; N=3 birds
bs	existence metabolism as measured by the difference between food energy and energy in egesta of birds maintained at constant body mass for periods of 3 or more days; N=NR; winter
bt	daily energy metabolism based on heart rate (30 C) and a linear regression equation for heart rate and volume of oxygen consumed; N=1; see citation for other metabolic estimates
bu	N=23; Condition=Breeding; February 19-July 8; agricultural lands (lat, 38 deg 32'N; long., 121 deg 47'W)
bv	N= 1 study area; Condition=Breeding; Mar.-Aug.; agricultural lands, woodlots, meadows
bw	N=NR; Condition=wintering; winter; southcentral area, agricultural lands; see citation for figure showing seasonal population size
bx	mean pair density in study area (194 km ²); N= 5 years; Condition=Breeding; rural and urban lands
by	number of individual kestrels counted annually on study site; N=4 years; Hanford Site, Benton and Franklin Counties
bz	number of nesting pairs counted annually on study site; N=4 years; Hanford Site, Benton and Franklin Counties
ca	density of nesting pairs; N=10 nests; spring; Jackson Hole
cb	average density; N=205 obs.; winter; agricultural lands; males predominant in areas with trees, females predominant in areas with few trees
cc	average density; N=135 obs.; winter; coastal and coast range areas; males predominant in areas with trees, females predominant in areas with few trees

cd	N= 1 study area; Sep.-Feb.; agricultural lands, woodlots, meadows
ce	mean density in study area; N=5 years; winter; rural and urban lands
cf	breeding population density; N=25 birds; January-June; agricultural lands
cg	wintering population size; N=NR; December-January; agricultural lands
ch	dorsal surface area; N=NR
ci	ventral surface area; N=NR
cj	mortality rate of banded individuals, 1946-65; N=6,544; throughout USA and Canada
ck	mortality rate estimate for banded individuals; N=118; throughout USA and Canada; kestrels banded as nestlings from 1925 to 1945
cl	average length of survival of banded individuals; N=1,017; throughout USA
cm	average annual mortality of banded individuals (over 6 years); N=NR; throughout USA; corresponds to life expectancy of 15 months for banded individuals
cn	hatching success; N=55 eggs (14 clutches); spring; Berks county
co	average diameter of territories; N=16; Condition=wintering; winter; southcentral area, agricultural lands
cp	N=5; winter; Hollister Basin (elev. 48-210 m)
cq	N=5; winter; Hollister Basin (elev. 48-210 m)
cr	observed range of movement during nesting, based on observations of individual or pair movements; N=11 birds; spring; Jackson Hole
cs	observed range of movement, based on observations of individuals; N=8 birds; winter; Superior Township
ct	median fledging date; N=27 pairs; Great Basin (elev. 1260-2340 m)
cu	median hatching date; N=27 pairs; Great Basin (elev. 1260-2340 m)
cv	dates of earliest and latest hatching; N=11 nests; spring; Jackson Hole
cw	median date of laying; N=27 pairs; Great Basin (elev. 1260-2340 m)
cx	period of laying; N=NR; Condition=breeding; spring; see citation for regional differences in date of laying initiation
cy	entire laying period; N=NR; Sagehen Creek Field Station
cz	period of fall migration of individuals fledged in northern CA, or at high elevations; N=NR; throughout state
da	period of spring migration of individuals fledged in northern CA, or at high elevations; N=NR; throughout state
db	time period of fall migration; N=276; fall; Pt. Diablo, Marin Headlands; see citation for monthly count data
dc	N=NR; Sagehen Creek Field Station
dd	period of molt; N=NR; throughout USA
de	period from earliest egg laying date to date of latest brood departure; N=11 nests; spring; Jackson Hole

References

1 Bloom, Peter H. and Stephen J. Hawks. 1983. Nest box use and reproductive biology of the American kestrel in Lassen County, California. *J. Raptor Res.* 17(1):9-14.

2 Porter, Richard D. and Stanley N. Wiemeyer. 1972. Reproductive patterns in captive American kestrels (sparrow hawks). *Condor.* 74:46-53.

3 Balgooyen, Thomas G. 1976. Behavior and ecology of the American Kestrel in the Sierra Nevada of California. *Univ. Calif. Publ. Zool.* 103:.

4 Harden, Shari M. 1993. Fat content of American kestrels and sharp-shinned hawks estimated by total body electrical conductivity. Master's thesis. Logan, UT: Utah State University, Logan. 42 p.

5 Gessaman, J.A. and S.M. Harden. 1995. Fat content of American kestrels and sharp-shinned hawks estimated by total body electrical conductivity. *J. Raptor Res.* 29(1):56-57.

6 Gessaman, James A. 1979. Premigratory fat in the American kestrel. *Wilson Bull.* 91(4):625-626.

7 Roest, A.I. 1957. Notes on the American sparrow hawk. *Auk.* 74(1):1-19.

8 Bird, David M. and Robert G. Clark. 1983. Growth of body components in parent- and hand-reared captive kestrels. *J. Raptor Res.* 17(3):77-84.

9 Bloom, Peter H. 1973. Seasonal variation in body weight of sparrow hawks in California. *West. Bird Bander.* 48(2):17-19.

10 Henry, Charles J. 1972. An analysis of the population dynamics of selected avian species with special reference to changes during the modern pesticide era. Washington, D. C.: United States Department of the Interior, Fish and Wildlife Service, Wildlife Research Report No. 1. 23-28 p.

11 Toland, Brian R. and William H. Elder. 1987. Influence of nest-box placement and density on abundance and productivity of American kestrels in central Missouri. *Wilson Bull.* 99:712-717.

12 Heintzelman, Donald S. and Alexander C. Nagy. 1968. Clutch sizes, hatchability rates, and sex ratios of sparrow hawks in eastern Pennsylvania. *Wilson Bull.* 80(3):306-311.

13 Willoughby, Ernest J. and Tom J. Cade. Breeding behavior of the American kestrel (sparrow hawk). *Living Bird.* 3:75-96.

14 Craighead, John J. and Frank C. Craighead, Jr. 1969. *Hawks, Owls and Wildlife.* New York, N.Y.: Dover Publications, Inc. 443 p.

15 Craft, Randall A. and Karen P. Craft. 1996. Use of free ranging American kestrels and nest boxes for contaminant risk assessment sampling: a field application. *J. Raptor Res.* 30(4):207-212.

16 Jacobs, Eugene A. 1995. American kestrel reproduction and dispersal in central Wisconsin. *J. Raptor Res.* 29(2):135-137.

17 Rudolph, Seri Gale. 1980. The hunting behavior of American kestrels: Applications of foraging theory. Master's thesis. Davis, CA: University of California at Davis. 64 p.

18 Meyer, Ruthe Lashe and Thomas G. Balgooyen. 1987. A study and implications of habitat separation by sex of wintering American kestrels (Falco sparverius). J. Raptor Res. 6:107-123.

19 Collopy, Michael W. and James R. Koplin. 1983. Diet, capture success, and mode of hunting by female American kestrels in winter. Condor. 85:369-371.

20 Collopy, Michael W. 1973. Predatory efficiency of American kestrels wintering in northwestern California. J. Raptor Res. 7(2):25-31.

21 Hawkins, Cole Carlson. 1991. Food habits and abundance of raptors in an agricultural area of Tulare County, California. Master's thesis. Fresno, CA: California State University, Fresno. 102 p.

22 Rohrbaugh, Ronald W., Jr., and Richard H. Yahner. 1997. Effects of macrohabitat and microhabitat on nest-box use and nesting success of American kestrels. Wilson Bull. 109(3):410-423.

23 Vincenty, Joseph A., III. 1974. A study of factors affecting nesting raptor populations in urban areas, Sacramento County, California - 1974. Sacramento, CA: California Department of Fish and Game, Wildlife Management Branch Administrative Report No. 74-5. 1-21 p.

24 Fitzner, R.E. 1980. Impacts of a nuclear energy facility on raptorial birds, In: Richard P. Howard and James F. Grove, eds. Workshop on Raptors and Energy Developments: Proceedings of a Workshop; January 25-26, 1980; Boise, Idaho. Boise, ID: U.S. Fish and Wildlife Service, p 9-33.

25 Barrett, Gary W. and Mackey, Carl V. 1975. Prey selection and caloric ingestion rate of captive American kestrels. Wilson Bull. 87(4):514-519.

26 Wing, Leonard and Anne H. Wing. 1939. Food consumption of a sparrow hawk. Condor. 41:168-170.

27 Rudolph, Seri G. 1982. Foraging strategies of American kestrels during breeding. Ecology. 63:1268-1276.

28 Enderson, James Harris. 1960. A population study of the sparrow hawk in east-central Illinois. Wilson Bull. 72(3):222-231.

29 Daniels, Kathleen and G.E. Duke. 1980. Factors influencing respiratory rates in four species of raptors. Comp. Biochem. Physiol., A , Comp. Physiol. 66:703-706.

30 Kuyt, E. 1972. Longevity in a captive sparrow hawk. Blue Jay. 30:197.

31 Clapp, Roger B., M. Kathleen Klimkiewicz and John H. Kennard. 1982. Longevity records of North American birds: Gaviidae through Alcidae. J. Field Ornithol. 53(2):81-124.

32 Keran, Doug. 1981. The incidence of man-caused and natural mortalities to raptors. J. Raptor Res. 15(4):108-112.

33 Gatehouse, S.N. and B.J. Markham. 1970. Respiratory metabolism of three species of raptors. Auk. 87:738-741.

34 Koplin, James R., Michael W. Collopy, Albert R. Bammann and Howard Levenson. 1980. Energetics of two wintering raptors. Auk. 97:795-806.

35 Gessaman, James A. 1980. An evaluation of heart rate as an indirect measure of daily energy metabolism of the American kestrel. Comp. Biochem. Physiol., A , Comp. Physiol. 65:273-289.

36 Toland, Brian R. 1987. The effect of vegetative cover on foraging strategies, hunting success and nesting distribution of American kestrels in central Missouri. J. Raptor Res. 21:14-20.

37 Mills, G. Scott. 1975. A winter population study of the American kestrel in central Ohio. Wilson Bull. 87:241-247.

38 Koplin, James R. 1973. Differential habitat use by sexes of American kestrels wintering in northern California. J. Raptor Res. 7(2):39-42.

39 Hayes, Stephen R. and James A. Gessaman. 1982. Prediction of raptor resting metabolism: comparison of measured values with statistical and biophysical estimates. J. Therm. Biol. 7:.

40 Bloom, Peter H. 1985. Raptor movements in California, In: Michael Harwood, eds. Hawk Migration Association of North America. Proceedings of Hawk Migration Conference IV. Medford, MA: p 313-323.

41 Binford, Laurence C. 1979. Fall migration of diurnal raptors at Pt. Diablo, California. West. Birds. 10(1):1-16.

Cal/EPA, OEHHA and the University of California Regents are not responsible for damages of any kind resulting from the use of or reliance on information in this report. Users are encouraged to consult the original data. Updated: February 1999.